arXiv:cond-mat/9912066v1 [cond-mat.str-el] 4 Dec 1999

Magnetization Jump in the Spin-$\frac{1}{2}$ XXZ Chain with Next-Nearest-Neighbour Coupling

Shunsaku Hirata*

Department of Physics, Kyushu University, Fukuoka 812-8581

(Received February 1, 2008)

The zero-temperature magnetization process of a one-dimensional $S = \frac{1}{2}$ spin system with next-nearest-neighbour coupling α and anisotropy δ is studied. There is a region in the parameter space where the magnetization has a jump at certain external field. The boundary of this region is determined. The two cases are distinguished according as $\alpha < \frac{1}{4}$ or $\alpha > \frac{1}{4}$. In the latter case, the energy dispersion of the low-lying excitations exhibits a double minimum structure.

KEYWORDS: quantum spin chain, magnetization jump, two magnon bound state, frustration

The magnetization process is one of the interesting topics in the study of low-dimensional quantum spin systems. In particular, the one-dimensional systems at low-temperature attracted a great deal of theoretical and experimental interest because of pronounced effect induced by quantum fluctuation in those systems. They exhibit various nontrivial behaviours, *e.g.*, the middle-field cusp singularity,[1] the magnetization plateau[2,3] and the second-order spin-flop transition.[4] What we concern in the present study is the magnetization jump similar to the metamagnetic transition in the classical Ising-like antiferromagnet, but it is caused by a different mechanism.

In this paper, we study a spin-$\frac{1}{2}$ XXZ chain with nearest-neighbour (NN) coupling J and next-nearest-neighbour (NNN) coupling $J\alpha$ in a magnetic field h. The model is described by the Hamiltonian

$$H = H_0 - h\sum_{i=1}^{L} S_i^z, \tag{1}$$

$$H_0 = J\sum_{i=1}^{L}(h_{i,i+1} + \alpha h_{i,i+2}), \tag{2}$$

$$h_{i,j} = S_i^x S_j^x + S_i^y S_j^y + \delta S_i^z S_j^z, \tag{3}$$

where $\boldsymbol{S}_i$'s represent the $S = \frac{1}{2}$ spin operators on the chain, and L is the total number of spins. The periodic boundary condition $\boldsymbol{S}_{i+L} = \boldsymbol{S}_i$ is assumed. Hereafter, we set $J > 0$ and take a unit so that $J = 1$. The model has been studied extensively. In the absence of external fields, it exhibits a variety of phenomena associated with competing interaction such as spontaneous dimerization[5,6,7] and incommensurate spin correlations.[8,9]

We shall concentrate on the region $0 \le \alpha \le \frac{1}{2}, -1 < \delta < 0$ where an anomalous behaviour in the magnetization process was reported:[10] In a region of parameter space, namely, $\delta_f(\alpha) < \delta < \delta_c(\alpha)$, there exists a critical field h_c at which the magnetization jumps to its maximum value $m_s = \frac{1}{2}$ (see Fig. 1). It should be noted that the model has competing NN and NNN exchange couplings in the easy-plane (XY-plane), while there is no competition in the z-axis although couplings are relatively weak. It is responsible for the existence of the magnetization jump: The system prefers to oder in the XY-plane but NNN coupling introduces frustration. Therefore, it becomes energetically preferable to order along the z-axis for sufficiently large α and $|\delta|$. For $\delta < \delta_f$, the zero-field ground state becomes the completely aligned (ferromagnetic) state with $m = \frac{1}{2}$.



Fig. 1. The zero-temperature magnetization curves for $\alpha = 0.25$ and $\delta = 0.0, -0.25, -0.5$. The thin lines show the finite-size results for $L = 18$ and 20. The magnetization (per site) m is normalized by the saturation magnetization $m_s = \frac{1}{2}$. The thick lines represent the estimated limiting ($L \to \infty$) curves. For $\delta = -0.25$ and -0.5, there exists a critical field h_c at which the magnetization jumps to the saturation value m_s.

The purpose of this paper is to determine the boundary $\delta = \delta_c(\alpha)$ of the region where the magnetization jump takes place.[11]

First, we discuss the condition that the magnetization jump occurs. For the magnetization curve to be smooth, $\epsilon_0(m)$, the lowest energy eigenvalue (per site) for the

---

* E-mail: hira4scp@mbox.nc.kyushu-u.ac.jp



Fig. 2. (a) Schematic curve of $\epsilon_0(m)$ in the case when the magnetization jump occurs. There is a region in which ϵ_0'' is negative. The line with slope h_c is also shown where h_c is a critical field. (b) Magnetization curve corresponding to ϵ_0 of (a).



Fig. 3. (a) The m-dependence of ϵ_0 and (b) magnetization curves of the $S = \frac{1}{2}$ XXZ chain with NNN interaction in the case when $\delta_f \le \delta \le \delta_c$. The dotted line indicates the critical magnetization $m_c(\delta)$.

Hamiltonian H_0 in the fixed $m = \sum S_i^z / L$ sector, must be monotonically increasing in m and convex for $0 < m < m_s = \frac{1}{2}$;

$$\epsilon_0''(m) = \frac{d^2 \epsilon_0(m)}{dm^2} > 0 \ . \tag{4}$$

On the other hand, if there exists a region where $\epsilon_0'' \le 0$, then the magnetization jumps over that region as schematically shown in Figs. 2(a) and 2(b). Thus, the presence of such a region as $\epsilon_0'' \le 0$ means that the magnetization jump occurs. The quantity $\epsilon_0''(m)$ can be estimated from

$$\epsilon_0''(m) = \lim_{L \to \infty} L \times \Delta(m), \tag{5}$$

where

$$\Delta(m) = E\left(m + \frac{1}{L}\right) + E\left(m - \frac{1}{L}\right) - 2E(m).$$

In the case of our model, the problem of determining the boundary curve $\delta = \delta_c(\alpha)$ is simplified. The results of the exact diagonalization suggest that the critical magnetization m_c increases as δ is increased and reaches m_s at $\delta = \delta_c$. It indicates that

$$\epsilon_0''|_{m=\frac{1}{2}-0} \le 0 \tag{6}$$

at least for $\delta \lesssim \delta_c$ [see Figs. 1, 3(a) and 3(b)]. The boundary δ_c is a line where $\epsilon_0''|_{m=\frac{1}{2}-0}$ changes sign. Therefore, we have merely to consider up to two-spin deviations from the completely aligned state for our purpose. We put $\Delta^{(1)} = \Delta(\frac{1}{2} - \frac{1}{L})$;

$$\Delta^{(1)} = E_0^{(2)} - E_0^{(0)} - 2\left[E_0^{(1)} - E_0^{(0)}\right], \tag{7}$$

where $E_0^{(0)}$ denotes the energy of the ferromagnetic state and $E_0^{(n)}$ the lowest energy of the states with n-spin deviations from the ferromagnetic state. For $\delta > \delta_f$, the ferromagnetic state is not a ground state of the Hamiltonian H_0. However, we shall call those n-deviation states as n-magnon states although they are not the excitations from the ground state.

We now obtain the energy spectrum of one- and two-magnon states. We rewrite the Hamiltonian (2) in term of fermion creation and annihilation operators with momentum k, a_k^+ and a_k, via Jordan-Wigner transformation. The Hamiltonian becomes

$$\begin{aligned} H = {} & E_0^{(0)} + \sum_k \omega_q a_q^+ a_q \\ & - \frac{1}{L} \sum_{1,2,3,4} \delta_{1+2,3+4} \Gamma_{1,2,3,4} a_1^+ a_2^+ a_3 a_4, \end{aligned} \tag{8}$$

where

$$\left\{ \begin{array}{rcl} E_0^{(0)} & = & \frac{1}{4}(1+\alpha)\delta L \\ \omega_q & = & \cos q - \delta + \alpha(\cos 2q - \delta) \\ \Gamma_{1,2,3,4} & = & \delta\cos(q_2 - q_4) + \alpha\delta\cos\left(2(q_2 - q_4)\right) \\ & & +2\alpha\cos(2q_1 + q_2 - q_3) \end{array} \right. ,$$

and we have used shorthand notations $1,2,3,4$ for q_1, q_2, q_3, q_4. In eq. (8), $E_0^{(0)}$ and ω_q represent the energy of the ferromagnetic state $|0\rangle$ and the excitation



Fig. 4. Energy spectrum of one-magnon states. For $\alpha > \frac{1}{4}$, two minima appears at $q = \pm q_0$ where $q_0 = \arccos(-\frac{1}{4\alpha})$.

spectrum of the one-magnon state with momentum q, respectively. For $\alpha \leq \frac{1}{4}$, ω_q has a single minimum at $q = \pi$, while, for $\alpha > \frac{1}{4}$, the minimum shifts from $q = \pi$ and two minima appears at $q = \pm q_0$ where $q_0 = \arccos(-\frac{1}{4\alpha})$ (Fig. 4).

The total wave number Q of two magnons to be conserved, thus we write two-magnon state as

$$|Q\rangle = \sum_{q>0} f_Q(q) a^+_{q_1} a^+_{q_2} |0\rangle, \tag{9}$$

where

$$Q = q_1 + q_2 \ , \ q = \frac{1}{2}(q_1 - q_2) \ .$$

Let us solve the eigenvalue equation

$$H|Q\rangle = E|Q\rangle, \tag{10}$$

where E is the energy eigenvalue of the two-magnon state. Taking the inner product of both side of eq. (10) with every basis state $\langle 0|a_{q_1} a_{q_2}$, we obtain the following equation:

$$[E - \gamma_{Q,q}] f_Q(q) = \frac{4}{L} \sum_{\kappa>0} f_Q(\kappa)[\chi_Q \sin q \sin \kappa + \alpha\delta \sin 2q \sin 2\kappa], \tag{11}$$

where

$$\left\{ \begin{array}{lcl} \gamma_{Q,q} & = & E_0^{(0)} + \omega_{q_1} + \omega_{q_2} \\ \chi_Q & = & \delta + 2\alpha \cos Q \end{array} \right. .$$

Here, $\gamma_{Q,q}$ represents the energy of the free two-magnons with total wave number $Q = q_1 + q_2$, and forms a continuum for each value of Q.

We are interested in whether there exists a solution of eq. (11) which gives $\Delta^{(1)} < 0$. Hence, it is sufficient to look for the solution satisfies $E < \gamma_{Q,q}$, i.e., the two-magnon bound states below the continuum [cf. eq. (7)]. Then, the eigenvalue equation reduces to (see, for instance, ref. 12)

$$D_Q(E) = \left| \begin{array}{cc} 1 - 2\chi_Q I_Q^{2,0} & -2\alpha\delta I_Q^{1,1} \\ -2\chi_Q I_Q^{1,1} & 1 - 2\alpha\delta I_Q^{0,2} \end{array} \right| = 0 \ , \tag{12}$$



Fig. 5. The results of $\Delta^{(1)}$ for $\alpha = 0.25$ and $\alpha = 0.3$. There is a point δ_c where $\Delta^{(1)}$ vanishes, which is a boundary of the region where the magnetization jump occurs.

where

$$\begin{aligned} I_Q^{l,m}(E) &= \frac{2}{L} \sum_{q>0} \frac{\sin^l q \sin^m 2q}{E - \gamma_{Q,q}} \\ &\xrightarrow{L\to\infty} \frac{1}{\pi} \int_0^\pi dq \frac{\sin^l q \sin^m 2q}{E - \gamma_{Q,q}}. \end{aligned} \tag{13}$$

It is the sufficient and necessary condition for the bound state lying outside of the continuum to exist; i.e., such bound states exists if and only if eq. (12) folds. The explicit solution of (12) is found analytically for several values of α, δ and Q. It is investigated numerically for general case.

Figure 5 shows the results of $\Delta^{(1)}$ for $\alpha = 0.25$ and 0.3. It can be seen that there exists a point δ_c at which $\Delta^{(1)}$ vanishes for each value of α and that $\Delta^{(1)}$ is negative for $\delta < \delta_c$ which suggests the magnetization jump to occur.[15)] In Fig. 6, the result is compared with the one obtained from the diagonalization of the finite chain Hamiltonian for rather large systems (L = 20-156) at $\alpha = 0.25, \delta = -0.25$. Note that there is a weak oscillatory behaviour in the finite-size results. For $\alpha > \frac{1}{4}$, there appears rapid oscillation, thus it becomes difficult to estimate a limiting value of $\Delta^{(1)}$ from the finite chain calculation.

In Fig. 7, the curve $\delta_c(\alpha)$ is shown in the α-δ plane. The magnetization jump occurs in the region enclosed by the two curves δ_c and δ_f.[16)] The ground-state phases in the absence of external fields are also shown;[17)] there are three phases, namely, the spin fluid phase which is characterized by the gapless excitation and critical behaviour, the spontaneously dimerized phase and the ferromagnetic phase. From Fig. 7, it can be clearly seen that the curve δ_c has a kink at $\alpha = \frac{1}{4}$. It results from the fact that the dispersion curve of the one-magnon states changes its shape at $\alpha = \frac{1}{4}$: For $\alpha < \frac{1}{4}$, it has a single minimum at $q = \pi$, while, for $\alpha > \frac{1}{4}$, two minima appears. The lowest lying state should consist of bound pairs of magnons with $q \simeq \pi$ for the former case and with $\pm q_0$ for the latter case.



Fig. 6. Comparison with the result from the numerical diagonalization. Open circles denotes the finite chain results for $L = 20$-156. The result obtained from eq. (12) is $\Delta^{(1)} = -1.43 \times 10^{-5}$.



Fig. 7. The phase diagram in the α-δ plane. The magnetization jump occurs in the region enclosed by the two curves δ_c (○) and δ_f (●). The zero-field ground-state phase diagram is also shown; the boundary of the ferromagnetic phase δ_f and the dimer-spin fluid phase boundary (dashed line).

To summarize, we determined the boundary of the region in which the magnetization jump occurs in the $S = \frac{1}{2}$ XXZ chain with easy-plane anisotropy and NNN interaction in the case of $0 < \alpha < \frac{1}{2}$. It was shown that the α-dependence of the boundary curve δ_c is qualitatively different according as $\alpha < \frac{1}{4}$ or $\alpha > \frac{1}{4}$. For $\alpha > \frac{1}{4}$, the region that the magnetization jump occurs shrinks with increasing α in contrast to the case of $\alpha < \frac{1}{4}$. It reflects the fact that the energy dispersion curve of the low-lying excitation develops the double minimum structure beyond the line $\alpha = \frac{1}{4}$.

One might be interested in how the system is in the case when the external field h is close to the critical field h_c. One of the possible situation is that there are clusters of up-spins separated by fluid-like domains.

The author would like to thank K. Nomura for valuable discussion and comments.

---

[1] K. Okunishi, Y. Hieida and Y. Akutsu: Phys. Rev. B **60** (1999) R6953.
[2] K. Hida: J. Phys. Soc. Jpn. **63** (1994) 2359.
[3] K. Okamoto: Solid State Commun. **98** (1995) 245.
[4] T. Sakai and M. Takahashi: Phys. Rev. B **60** (1999) 7295.
[5] F. D. M. Haldane: Phys. Rev. B **25** (1982) 4925.
[6] K. Okamoto and K. Nomura: Phys. Lett. A **169** (1992) 433.
[7] K. Nomura and K. Okamoto: J. Phys. A **27** (1994) 5773.
[8] T. Tonegawa and I. Harada: J. Phys. Soc. Jpn. **56** (1987) 2153.
[9] S. R. White and I. Affleck: Phys. Rev. B **54** (1996) 9862.
[10] C. Gerhardt, K.-H. Mütter and H. Kröger: Phys. Rev. B **57** (1998) 11504.
[11] In ref. 10, Gerhardt *et al.* determined the boundary δ_c by use of the numerical diagonalization of finite chain Hamiltonian up to $L = 50$. However, their result suffers from finite size effects and somewhat suspicious (compare Fig. 6 in this work with FIG. 3 of ref. 10).
[12] I. Ono, S. Mikado and T. Oguchi: J. Phys. Soc. Jpn. **30** (1971) 358.
[13] The integrand of (13) has, however, a sharp peak which becomes singular as E approaches to the lower edge of the scattering continuum $\gamma_{Q,q}$. Thus, for numerical quadrature we use the double exponential formula,[14] which is one of the most optimal and suitable numerical quadrature formula for improper integrals.
[14] H. Takahashi and M. Mori: Publ. RIMS, Kyoto Univ. **9** (1974) 721.
[15] To discuss thermodynamic limit, we must take into account the finite size correction: If $\Delta^{(1)}$ behaves as $\Delta^{(1)} = \Delta^{(1)}_{\infty} + c_1 L^{-1} + O(L^{-2})$ and if $\Delta^{(1)}_{\infty} = 0$, then it is the coefficient of the correction term c_1 that determine the sign of ϵ''_0 in thermodynamic limit. We are unable to estimate the finite size correction analytically. However, it seems that there is no significant difference in δ_c.
[16] The boundary of the ferromagnetic phase δ_f was determined from the level crossing of the ferromagnetic state and the lowest eigenstate with $m = 0$ (see ref. 17 for details).
[17] S. Hirata and K. Nomura: cond-mat/9909290.